

A Remark on the Three Dimensional Baroclinic Quasigeostrophic Dynamics *

Jinqiao Duan

Department of Mathematical Sciences,
Clemson University, Clemson, South Carolina 29634, USA.
E-mail: duan@math.clemson.edu, Fax: (864)656-5230.

January 12, 1998

Abstract

We consider time-periodic patterns of the dissipative three dimensional baroclinic quasigeostrophic model in spherical coordinates, under time-dependent forcing. We show that when the forcing is time-periodic and the spatial square-integral of the forcing is bounded in time, the model has time-periodic solutions.

Keywords— quasigeostrophic fluid model, dissipative dynamics, time-periodic motion, nonlinear analysis.

Short running title:

Three Dimensional Quasigeostrophic Dynamics

*This research was supported by the National Science Foundation Grant DMS-9704345 and a Clemson University Research Grant. This research was performed while the author was visiting the Isaac Newton Institute for Mathematical Sciences at Cambridge University, England.

1 Introduction

The three dimensional baroclinic quasigeostrophic model is an approximation of the rotating Euler (inviscid case) or Navier-Stokes (viscous case) equations in the limit of zero Rossby number, i.e., at asymptotically high rotation rate; see, for example, [1], [2], [3], [4], [5], [6], and [7].

The well-posedness for the inviscid three dimensional baroclinic quasigeostrophic model was studied in, for example, [8] and [4], and for the viscous three dimensional baroclinic quasigeostrophic model in [9]. The inviscid baroclinic quasigeostrophic model is a Hamiltonian system ([10]). The work mentioned above and most other research about the baroclinic quasigeostrophic model has been on the Cartesian, β -plane version of this model.

Wang ([11]) discussed global attractors for the viscous three dimensional baroclinic quasigeostrophic model in spherical coordinates, under time-independent forcing.

In this paper, we consider the same viscous three dimensional baroclinic quasigeostrophic model, but under time-dependent forcing. We show that when the forcing is time-periodic and when the spatial square-integral of the forcing is bounded in time, the forced viscous three dimensional quasigeostrophic model has time-periodic solutions. We use a topological technique from nonlinear global analysis ([12]).

2 Dissipative Dynamics and Time-Periodic Motion

We consider the three dimensional baroclinic quasigeostrophic equation, in nondimensional form, for the atmospheric dynamics in spherical coordinates ϕ, θ, ζ as in [11]

$$(A\psi)_t + J(\psi, Ro A\psi + 2\cos\theta) - \frac{1}{Re}A^2\psi = f(\phi, \theta, \zeta, t), \quad (1)$$

where $\psi(\phi, \theta, \zeta, t)$ is the stream function, $f(\phi, \theta, \zeta, t)$ is the time-dependent forcing such as external source of heating, and Ro, Re are the Rossby number, Reynolds number, respectively. The space domain for the equation is $D = S^2 \times (\zeta_0, 1)$ with S^2 the two dimensional unit sphere and ζ_0 fixed: $1 > \zeta_0 \geq 0$.

The operators appeared in the equation (1) are

$$\Delta = \frac{1}{\sin\theta} \left[\frac{\partial}{\partial\theta} (\sin\theta \frac{\partial}{\partial\theta}) + \frac{1}{\sin\theta} \frac{\partial^2}{\partial\phi^2} \right],$$

$$A = \Delta + \frac{\partial}{\partial\zeta} (N^2 \frac{\partial}{\partial\zeta}),$$

$$J(p, q) = \frac{\partial p}{\partial\theta} \frac{1}{\sin\theta} \frac{\partial q}{\partial\phi} - \frac{1}{\sin\theta} \frac{\partial p}{\partial\phi} \frac{\partial q}{\partial\theta},$$

where $N(\phi, \theta, \zeta) > 0$ is a known smooth function related to the stratification frequency.

The equation (1) is supplemented by the following boundary and initial conditions

$$\frac{\partial\psi}{\partial\zeta} = \frac{\partial(A\psi)}{\partial\zeta} = 0 \quad \text{if } \zeta = \zeta_0, \quad (2)$$

$$\frac{\partial\psi}{\partial\zeta} + a\psi = \frac{\partial(A\psi)}{\partial\zeta} + aA\psi = 0 \quad \text{if } \zeta = 1, \quad (3)$$

$$\psi|_{t=0} = \psi_0(\phi, \theta, \zeta), \quad (4)$$

where $a(\phi, \theta) > 0$ is a known smooth function related to heat transfer between the atmosphere and the earth, and ψ_0 is initial data.

Denote $L^2(D)$ as the space of square-integrable functions, with the standard norm $\|\cdot\|$. The problem (1), (2), (3), (4) is well-posed ([11]) in $L^2(D)$, with solution $\psi(\phi, \theta, \zeta, t)$ at least continuous in time t .

We address the issue of whether there are any time-periodic solutions in the nonlinear forced dissipative quasigeostrophic dynamics modeled by (1), (2), (3). To this end we further assume that the forcing $f(t) := f(\phi, \theta, \zeta, t)$ is periodic in time with period $T > 0$, and the spatial square-integral of the forcing $\|f(t)\|$ is bounded in time, i.e., $\|f(t)\|$ is bounded by a time-independent constant. Then we can follow [11] exactly to show that there is a bounded absorbing set in $L^2(D)$ (we omit this part). That is, all solutions ψ enter a bounded set $\{\psi : \|\psi\| \leq C(Re, Ro)\}$ as time goes to infinity. The system (1), (2), (3) is therefore a dissipative system as defined in [12] (also [13] or [14]).

We now recall a result from [12], page 235, that a T -time-periodic nonautonomous dissipative dynamical system in a Banach space has at least one T -time-periodic solution. This result follows from a Leray-Schauder topological degree argument and the Browder's principle ([12]). So the system (1),

(2), (3) has at least one T -time-periodic solution starting from some initial data ψ_0 . Therefore we have the following result.

Theorem 1 *Assume that the forcing $f(\phi, \theta, \zeta, t)$ is time-periodic with period $T > 0$, and its spatial square-integral with respect to ϕ, θ, ζ is bounded in time. Then the forced viscous three dimensional baroclinic quasigeostrophic model*

$$(A\psi)_t + J(\psi, Ro A\psi + 2\cos\theta) - \frac{1}{Re} A^2\psi = f(\phi, \theta, \zeta, t), \quad (5)$$

$$\frac{\partial\psi}{\partial\zeta} = \frac{\partial(A\psi)}{\partial\zeta} = 0 \quad \text{if } \zeta = \zeta_0, \quad (6)$$

$$\frac{\partial\psi}{\partial\zeta} + a\psi = \frac{\partial(A\psi)}{\partial\zeta} + aA\psi = 0 \quad \text{if } \zeta = 1, \quad (7)$$

$$(8)$$

has at least one time-periodic solution with period $T > 0$, for some square-integrable initial data ψ_0 .

3 Discussions

Forced coherent structures in the two dimensional baroclinic model were studied in [15]. A wind forced two dimensional baroclinic model was also used in the study of multiple geophysical equilibria (e. g., [16]). In general, it is very difficult to show existence of periodic coherent structures in spatially extended physical systems. In this paper, we have shown that the three dimensional time-periodic quasigeostrophic patterns may form due to time-periodic forcing such as external source of heating.

References

- [1] J. G. Charney, The dynamics of long waves in a baroclinic westerly current, *J. Meteorol.* (1947), 135-163.
- [2] J. Pedlosky, *Geophysical Fluid Dynamics*, Springer-Verlag, 2nd edition, 1987.
- [3] A. E. Gill, *Atmosphere-Ocean Dynamics*, Academic Press, New York, 1982.

- [4] A. J. Bourgeois and J. T. Beale, Validity of the quasigeostrophic model for large-scale flow in the atmosphere and ocean, *SIAM J. Math. Anal.* **25** (1994), 1023-1068.
- [5] P. F. Embid and A. J. Majda, Averaging over fast gravity waves for geophysical flows with arbitrary potential vorticity, *Comm. PDEs* **21** (1996), 619-658.
- [6] B. Desjardins and E. Grenier, Derivation of quasigeostrophic potential vorticity equations, preprint, 1997.
- [7] B. Desjardins and E. Grenier, On the homogeneous model of wind driven ocean circulation, preprint, 1997.
- [8] J. A. Dutton, The nonlinear quasi-geostrophic equation: Existence and uniqueness of solutions on a bounded domain, *J. Atmos. Sci.* **31** (1974), 422-433.
- [9] A. F. Bennett and P. E. Kloeden, The dissipative quasigeostrophic equations, *Mathematika* **28** (1981), 265-285.
- [10] D. D. Holm, Hamiltonian formulation of the baroclinic quasigeostrophic fluid equations, *Phys. Fluids* **29** (1986), 7-8.
- [11] S. Wang, Attractors for the 3D baroclinic quasi-geostrophic equations of large-scale atmosphere, *J. Math. Anal. Appl.* **165** (1992) 266-283.
- [12] M. A. Krasnoselskii and P. P. Zabreiko, *Geometrical Methods of Nonlinear Analysis*, Springer-Verlag, New York, 1984.
- [13] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, Springer-Verlag, New York, 1988.
- [14] J. K. Hale, *Asymptotic Behavior of Dissipative Systems*, American Math. Soc., 1988.
- [15] R. T. Pierrehumbert and P. Malguzzi, Forced coherent structures and local multiple equilibria in a barotropic atmosphere, *J. Atmos. Sci.* **41** (1984), 246-257.
- [16] P. Cessi and G. R. Ierley, Symmetry-breaking multiple equilibria in quasigeostrophic, wind-driven flows, *J. Phys. Oceanography*, **25** (1995), 1196-1205.